

Journal: European Heart Journal – Cardiovascular Imaging

Article id: jet063




Article title: Prodromal angina is associated with myocardial salvage in acute ST-segment elevation myocardial infarction

First Author: Pier Giorgio Masci

Corr. Author: Pier Giorgio Masci

AUTHOR QUERIES - TO BE ANSWERED BY THE CORRESPONDING AUTHOR

The following queries have arisen during the typesetting of your manuscript. Please answer these queries by marking the required corrections at the appropriate point in the text. Failure to do so could result in delayed publication.

Query No.	Nature of Query	Author's Response
Q1	Please check that all names have been spelled correctly and appear in the correct order. Please also check that all initials are present. Please check that the author surnames (family name) have been correctly identified by a pink background. If this is incorrect, please identify the full surname of the relevant authors. Occasionally, the distinction between surnames and forenames can be ambiguous, and this is to ensure that the authors' full surnames and forenames are tagged correctly, for accurate indexing online. Please also check all author affiliations.	
Q2	Figures have been placed as close as possible to their first mention in the text. Please check that the figures are accurately placed in the text, that the images are correct, and that they have the correct caption and citation.	
Q3	Please check the suggested short title for the article.	

MAKING CORRECTIONS TO YOUR PROOF

These instructions show you how to mark changes or add notes to the document using the Adobe Acrobat Professional version 7.0 (or onwards) or Adobe Reader 8 (or onwards). To check what version you are using go to **Help** then **About**. The latest version of Adobe Reader is available for free from get.adobe.com/reader.

For additional help please use the **Help** function or, if you have Adobe Acrobat Professional 7.0 (or onwards), go to http://www.adobe.com/education/pdf/acrobat_curriculum7/acrobat7_lesson04.pdf

Displaying the toolbars

Adobe Reader 8: Select Tools, Comments & Markup, Show Comments and Markup Toolbar. If this option is not available, please let me know so that I can enable it for you.



Acrobat Professional 7: Select Tools, Commenting, Show Commenting Toolbar.




Adobe Reader 10: To edit the galley proofs, use the Comment Toolbar (Sticky Note and Highlight Text).



Using Text Edits

This is the quickest, simplest and easiest method both to make corrections, and for your corrections to be transferred and checked.




1. Click **Text Edits**
2. Select the text to be annotated or place your cursor at the insertion point.
3. Click the **Text Edits** drop down arrow and select the required action.

You can also right click on selected text for a range of commenting options.


Pop up Notes

With *Text Edits* and other markup, it is possible to add notes. In some cases (e.g. inserting or replacing text), a pop-up note is displayed automatically.



To **display** the pop-up note for other markup, right click on the annotation on the document and selecting **Open Pop-Up Note**.

To **move** a note, click and drag on the title area.



To **resize** of the note, click and drag on the bottom right corner.



To **close** the note, click on the cross in the top right hand corner.



To **delete** an edit, right click on it and select **Delete**. The edit and associated note will be removed.

SAVING COMMENTS

In order to save your comments and notes, you need to save the file (**File, Save**) when you close the document. A full list of the comments and edits you have made can be viewed by clicking on the Comments tab in the bottom-left-hand corner of the PDF.



Prodromal angina is associated with myocardial salvage in acute ST-segment elevation myocardial infarction

Pier Giorgio Masci^{1*}, Daniele Andreini², Marco Francone³, Erika Bertella², Laura De Luca⁴, Michele Cocceani¹, Saima Mushtaq², Massimiliano Mariani¹, Iacopo Carbone³, Gianluca Pontone², Luciano Agati⁴, Jan Bogaert⁵, and Massimo Lombardi¹

¹Fondazione CNR/Regione Toscana 'G. Monasterio', Via Moruzzi 1, 56124 Pisa, Italy; ²Centro Cardiologico Monzino, IRCCS, Milano, Italy; ³Radiology Department, 'La Sapienza' University, Rome, Italy; ⁴Cardiology Department, 'La Sapienza' University, Rome, Italy; and ⁵Radiology Department, University Hospitals, Leuven, Belgium

Received 29 November 2012; accepted after revision 27 March 2013

Aims

Previous studies have shown that prodromal angina (PA) occurs frequently in acute myocardial infarction (MI) patients. However, the potential benefits of PA on ischaemic myocardial damage remain unknown.

Methods and results

One-hundred and fifty-four patients with acute ST-segment elevation MI successfully treated with primary percutaneous coronary intervention (PPCI) were prospectively evaluated for new-onset PA in the week preceding infarction and other factors known to influence myocardial salvage. Cardiovascular magnetic resonance was performed 8 ± 3 days after MI for the assessment of area-at-risk (AAR), MI size, myocardial haemorrhage (MH), microvascular obstruction (MO), and myocardial salvage index (MSI).

Patients with PA ($n = 60$) compared with those without PA ($n = 94$) showed similar AAR but significantly smaller MI size leading to larger MSI (0.53 ± 0.27 vs. 0.32 ± 0.26 , $P < 0.001$). Additionally, patients with PA had lower incidence of MH (18 vs. 33%) and MO (22 vs. 46%) than non-PA patients (both $P < 0.05$). At univariate analysis, higher MSI was associated with new-onset PA, lower myocardial oxygen consumption before PPCI, shorter time-to-PPCI, and higher post-procedural TIMI flow-grade. Neither collateral circulation nor medications administered before PPCI were associated to MSI. After correction for other covariates by multivariate analysis, new-onset PA remained significantly associated with MSI (β -value: 0.352, $P < 0.001$).

Conclusion

In acute MI patients, new-onset PA is associated with higher MSI independent of others factors known to influence jeopardized myocardium, as well as with less microvascular damage.

Keywords

Prodromal angina • Myocardial infarction • Cardioprotection

Introduction

The pathophysiological basis of ischaemic myocardial damage has been deeply investigated thoroughly in animal models of reperfused and non-reperfused myocardial infarction (MI), yet remains poorly understood in humans.^{1,2} This apparent paradox may be explained considering that: (i) in the clinical scenario many variables, which are difficult to account for, may influence ischaemic myocardial damage; (ii) the complexities inherent in the study of

jeopardized myocardium in humans. Experimentally, most of the variation in infarct size is related to area-at-risk (AAR) and reperfusion status, and to a lesser extent to collaterals, haemodynamic determinants of myocardial metabolic demand, and medications.¹ Moreover, brief episodes of myocardial ischaemia before sustained ischaemia reduces MI size, a phenomenon known as ischaemic preconditioning (IPC).³ In humans, acute MI is often preceded by prodromal angina (PA), and previous clinical studies suggested that PA may limit MI size through various mechanisms, including collaterals

* Corresponding author. Tel: +39 050 315 2824; Fax: +39 050 315 2166, Email: masci@ftgm.it

Published on behalf of the European Society of Cardiology. All rights reserved. © The Author 2013. For permissions please email: journals.permissions@oup.com

recruitment,⁴ reperfusion facilitation,⁵ and IPC.^{6–11} However, these studies were conducted in highly selected populations and limited by the retrospective nature. Therefore, whether PA confers cardioprotection in patients with acute MI is still a matter of debate.

Cardiovascular magnetic resonance (CMR) is a valuable tool to investigate this issue because it allows an accurate and comprehensive assessment of ischaemic myocardial damage in the early post-infarction period.^{12,13} Based on these premises, we prospectively studied a cohort of acute MI patients with the aim of examining the influence of PA on myocardial salvage using CMR.

Methods

Study population

Between January 2011 and January 2012, 212 consecutive acute ST-segment elevation MI patients [95 at Fondazione 'G. Monasterio'-Italy, Pisa & Massa (Centre A), 63 at La Sapienza University, Rome, Italy (Centre B) and 54 at Centro Cardiologico Monzino, Milan, Italy (Centre C)] were prospectively evaluated for study enrolment. CMR was performed 8 ± 3 days after MI. Inclusion criteria were: (i) chest pain suggestive of myocardial ischaemia lasting >30 min; (ii) ECG showing ST-segment elevation >0.1 mV in ≥ 2 limb leads or >0.2 mV in ≥ 2 contiguous precordial leads, or presumed new left bundle-branch block; (iii) successful treatment with primary percutaneous coronary intervention (PPCI) within 12 h from symptoms onset; (iv) ability to provide detailed clinical history. Exclusion criteria were prior MI or revascularization, atrial fibrillation, Killip class $>II$, renal failure [creatinine >2 mg/dL ($0.226 \mu\text{mol/dL}$)] and contraindications to CMR. The study complied with the Declaration of Helsinki, the Ethics Committee approved the research protocol, and informed consent was obtained from each subject.

Clinical variables

Within 5 days from admission, clinical history was recorded from each patient by one of the physicians participating to the study. Particular attention was paid to assess new-onset PA, defined as one or more episode of typical chest pain lasting ≤ 30 min at rest or on effort within 7 days from the infarction.^{5,6} The number of PA episodes and the time intervals between the first and last episode of PA and infarction were registered. Time-to-PPCI was defined as the interval time between the onset of continuous chest pain and the re-opening of the infarct-related artery by PPCI.

Blood pressure and heart rate values before PPCI were collected from clinical documents completed in the ambulance, emergency department, intensive coronary care unit, and catheterization laboratory. Mean rate pressure product (mRPP), which correlates with myocardial oxygen consumption before PPCI,¹⁴ was calculated by multiplying mean systolic blood pressure and mean heart rate. The use of nitrates, morphine, beta-blockers, and Gp IIb/IIIa before PPCI were also recorded.

Coronary angiography analysis

All coronary angiograms were analysed at Centre A by an experienced physician blinded to clinical and CMR data. The TIMI flow-grade of the infarcted-related artery was assessed before and after PPCI. The grade of epicardial collaterals to the infarcted-related artery was evaluated according to Rentrop *et al.*¹⁵; collateral circulation was defined as Rentrop grade ≥ 2 . Double- and triple-vessel coronary disease were

diagnosed when a $\geq 75\%$ stenosis in one or two vessels remote from the infarct-related artery, respectively. All patients received dual anti-platelet therapy upon admission.

CMR protocol

CMR studies were performed at Centre A with 1.5-T unit (CVI, GE-Healthcare, Milwaukee, USA), at Centre B with 1.5-T unit (Avanto Siemens, Erlangen, Germany), and at Centre C with 1.5-T unit (Discovery MR450, GE-Healthcare, Milwaukee, USA). All studies were performed using dedicated cardiac software, phased-array surface receiver coil, and electrocardiogram triggering. A similar CMR study protocol was followed in all centres (see Supplementary material online). Breath-hold steady-state free-precession cine CMR was performed in vertical and horizontal long-axis, and in short-axis orientations. In short-axis, both ventricles were completely encompassed by a stack of contiguous slices. Breath-hold black-blood T2-weighted short inversion-time inversion-recovery fast spin-echo (T2w-imaging) was performed in short-axis orientation for AAR quantification. Breath-hold contrast-enhanced segmented T1-weighted inversion-recovery gradient-echo sequence [late-gadolinium enhancement (LGE)-imaging] was used to quantify MI and microvascular obstruction (MO). 10–20 min after an intravenous bolus of 0.1 mmol/kg Gadolinium-BOPTA (Multihance-Bracco, Milan, Italy) or 0.2 mmol/kg Gadolinium-BOPTA Gadolinium-DOTA (Dotarem Guerbet Roissy, France) LGE-imaging was performed. Inversion time was individually adapted to nullify the signal of remote myocardium (usual range 220–300 ms).

Image analysis

All CMR studies were analysed off-line by the consensus of two experienced observers at Centre A using a workstation (Advantage Workstation, GE-Healthcare, Milwaukee, USA) with a dedicated software (MASS 6.1, Medis, Leiden, the Netherlands). Operators were unaware of clinical and angiographic data. Analysis was started by scoring T2w-imaging quality using a 4-grade score: (1) poor, (2) moderate, (3) good, and (4) excellent. Only exams with a score >2 were included. T2w-imaging was used to quantify AAR and visualize myocardial haemorrhage (MH). Myocardium with a signal intensity (SI) >2 SD above mean SI of remote non-infarcted myocardium was considered AAR. Attention was paid to exclude increased SI from the blood adjacent to endocardium ('slow flow'). MH was defined as a hypo-intense area in the core of AAR. When present, MH was included in hyper-intense myocardium for AAR calculation.¹² On post-contrast imaging, LGE was considered present if SI of the hyperenhanced myocardium was >5 SD above the mean SI of remote myocardium,¹⁶ whereas MO was defined as hypo-enhanced region within the infarcted myocardium. When present, MO was included in the hyper-intense myocardium for LGE quantification. Left-ventricular (LV) LGE and MO were expressed as absolute (grams) or relative (LV percentage) value. Infarct transmural was calculated as the ratio of mean thickness of hyperenhanced myocardium to mean thickness of the corresponding myocardial wall, multiplied by 100. Cine CMR was used to derive LV volumes, ejection-fraction, regional wall motion, and mass. For LV wall motion analysis, we used the 17-segment model proposed by AHA¹⁷ (segment 17 was excluded from analysis). Regional wall motion was scored for each segment from 1 to 5 (1: normal/mild hypokinesia; 2: moderate hypokinesia; 3: severe hypokinesia; 4: akinesia; 5: dyskinesia). Wall motion score index (WMSI) was determined as the sum of segmental scores divided by the number of segments.¹³

Statistical analysis

Continuous variables were expressed as mean \pm SD or median (25th–75th percentiles). Kolmogorov–Smirnov test was used to assess the normality of data distribution, and variables distributed non-normally were logarithmically transformed. Categorical variables were expressed as frequency with percentage. Student's independent t- or Mann–Whitney tests were used, as appropriate, to compare continuous variable differences. Comparisons between categorical variables were made with χ^2 test or Fisher's exact test, if an expected cell count was <5 . Correlations between continuous variables were assessed using Pearson's (r) coefficient or Spearman's rho, as appropriate. Univariate linear regression analysis was utilized to determine the association of variables with MSI. Variables known to influence MSI were tested as potential covariates.^{1–10} Multivariate linear regression analysis was used to evaluate the influence of covariates on MSI. Variables with a P -value <0.10 at univariate analysis were entered in the multivariable model as covariates. Statistical analysis was performed by SPSS software for Windows (15.0 release; SPSS, Chicago, IL, USA); all tests were two-tailed at 5% significance level.

Results

Study population

Thirty-five (18%) patients were excluded from the study because of inadequate T2w-imaging quality, thus leaving a total of 154 patients for the analysis (124 men, age 62 ± 11 years; Figure 1). Patients were dichotomized according to the presence ($n = 60$) or absence ($n = 94$) of new-onset PA. Baseline characteristics

are summarized in Table 1. Peak troponin I was lower in patients with than those without PA. Notably, mRPP before PPCI, TIMI flow-grade before and after PPCI and Rentrop-grade were similar in the two groups. Nitrate use before PPCI was more common in patients with PA.

Prodromal angina characteristics

The characteristics of PA are summarized in Table 2. Twenty-eight patients (47%) referred only one episode of PA, whereas the remaining 32 patients (53%) had repeated episodes in the week preceding infarction. Notably, 58 (97%) and 31 (52%) patients experienced the last episode of PA within 48 and 6 h from infarction, respectively. The two patients presenting PA > 48 h had one single episode of angina occurring 55 and 72 h before infarction.

Prodromal angina and ischaemic myocardial damage

CMR findings are summarized in Table 3. Patients with and without PA showed similar AAR although LGE extent was significantly lower in patients with PA, resulting in higher MSI. Infarct transmural-ity was also lower in patients with PA compared with those without PA (Figure 2). As a result, patients with PA had better regional and global LV systolic function than patients without PA. In addition, aborted MI (evidence of AAR on T2w-imaging but the absence of LGE on LGE-imaging) was more commonly observed in patients with than those without PA [8(13%) vs. 3(3%), $P = 0.017$]. Patients with PA also showed lower incidence of MH




Figure 1 Study protocol.

Table 1 Baseline patients characteristics

Characteristics	Presence of PA (n = 60)	Absence of PA (n = 94)	P-value
Age (years)	60 ± 12	63 ± 10	0.146
Male, n (%)	49 (82)	75 (80)	0.774
Cardiovascular risk factors, n (%)			
Hypertension	30 (50)	54 (58)	0.328
Diabetes	9 (15)	17 (18)	0.598
Hypercholesterolaemia	23 (38)	42 (45)	0.404
Family history CAD	22 (37)	39 (42)	0.516
Current smokers	32 (53)	43 (46)	0.391
Killip class II	2 (3)	7 (7)	0.299
Peak troponin I (ng/dL)	31 [10–77]	71 [22–115]	0.002
Time-to-PPCI (min)	245 ± 162	248 ± 166	0.908
Haemodynamic parameters before PPCI			
Mean systolic BP (mmHg)	138 ± 19	135 ± 17	0.232
Mean diastolic BP (mmHg)	82 ± 11	78 ± 10	0.004
Mean heart rate (bpm)	76 ± 13	75 ± 16	0.692
Mean RPP ([bpm] mmHg)	10 430 ± 2419	10 136 ± 2846	0.514
Medications before PPCI, n (%)			
Beta-blocker	7 (12)	9 (10)	0.659
Nitrate	29 (51)	30 (33)	0.030
Morphine	13 (22)	22 (24)	0.833
Gp IIb/IIIa	7 (11)	18 (19)	0.152
Catheterization laboratory data			
Infarct-related artery, n (%)			0.497
LAD	33 (55)	44 (46)	
LCx	8 (13)	12 (13)	
RCA	19 (32)	38 (41)	
Double-vessel disease, n (%)	20 (33)	28 (30)	0.643
Triple-vessel disease, n (%)	7 (12)	7 (7)	0.324
Use of Gp IIb/IIIa, n (%)	26 (44)	42 (45)	0.968
Thrombus aspiration n (%)	30 (50)	51 (54)	0.606
TIMI flow-grade pre-PPCI, n (%)			0.647
0/1	55 (92)	88 (94)	
2/3	5 (8)	6 (5)	
TIMI flow-grade post-PPCI, n (%)			0.320
0/1	2 (3)	1 (1)	
2/3	58 (97)	93 (99)	
Presence of collaterals, n (%)	5 (8)	12 (13)	0.392
Rentrop grade	0.23 ± 0.70	0.32 ± 0.78	0.489
Medications at discharge, n (%)			
ACE-i/ARBs	48 (84)	80 (87)	0.640
Beta-blocker	49 (86)	74 (80)	0.387
Statins	54 (95)	89 (97)	0.646
Spironolactone	10 (17)	21 (22)	0.440

ACE, angiotensin-converting enzyme; ARBs, angiotensin receptor blocker; BP, blood pressure; CAD, coronary artery disease; LAD, left-anterior descending coronary artery; LCx, left circumflex coronary artery; PA, prodromal angina; PPCI, primary percutaneous coronary intervention; RCA, right coronary artery; RPP, rate pressure product; TIMI, thrombolysis in myocardial infarction.

and MO. When including only patients with pre-PPCI TIMI flow-grade ≤ 1 and Rentrop-grade ≤ 1 ($n = 128$), MSI remained higher and infarct transmuralty was lower in patients with PA than those without PA (0.49 ± 0.26 vs. 0.34 ± 0.26 , $P = 0.003$ and 57 ± 32 vs. $72 \pm 30\%$, $P = 0.029$; respectively).

The influence of PA on jeopardized myocardium

In the overall population, MSI was 0.40 ± 0.28 and had a normal distribution (Kolmogorov–Smirnov: 1.041, $P = 0.229$). There was

Table 2 Characteristics of prodromal angina

Characteristics	Range	Median [25th–75th]
Number of episodes of PA	1–9	2 [1–3]
Duration of first PA episode (min)	2–30	15 [6–20]
Time from PA first to acute MI (h)	0.2–167	10 [4–34]
Duration of last PA last (min)	3–30	20 [10–25]
Time from PA last to acute MI (h)	0.2–72	6 [1–22]

PA, prodromal angina; MI, myocardial infarction.

Table 3 Cardiovascular magnetic resonance characteristics

Variables	Presence of PA (n = 60)	Absence of PA (n = 94)	P-value
Area-at-risk (g)	29 ± 17	30 ± 18	0.832
LGE extent (g)	10 [3–24]	18 [9–30]	0.009
LGE extent (% LV)	8 [3–19]	14 [8–23]	0.007
MSI	0.53 ± 0.27	0.32 ± 0.26	<0.001
MI transmural (%)	53 ± 33	74 ± 29	0.001
MH, n (%)	11 (18)	31 (33)	0.047
MO, n (%)	13 (22)	40 (43)	0.008
MO extent (g)	5 [4–7]	4 [2–8]	0.437
LV-EDV (mL)	141 ± 36	145 ± 36	0.515
LV-ESV (mL)	64 ± 29	71 ± 26	0.136
LV-SV (mL)	77 ± 19	74 ± 21	0.463
CI (mL/m ²)	2.69 ± 0.69	2.71 ± 0.74	0.848
LV-EF (%)	56 ± 10	52 ± 10	0.029
LV-mass (g)	135 ± 30	132 ± 37	0.529
LV-WMSI	1.502 ± 0.334	1.660 ± 0.405	0.030

CI, cardiac index; EDV, end-diastolic volume; EF, ejection fraction; ESV, end-systolic volume; LV, left ventricular; MI, myocardial infarction; MO, microvascular obstruction; MSI, myocardial salvage index; WMSI, wall motion score index.

a weak inverse relationship between MSI and time-to-PPCI ($r = -0.173$, $P = 0.031$). Dichotomizing patients on the basis of PA, MSI remained inversely related to time-to-PPCI in patients without PA ($r = -0.291$, $P = 0.004$) but not in those with PA ($r = -0.025$, $P = 0.850$). Similarly, in the overall population, MSI was inversely related to pre-PPCI mRPP ($r = -0.191$, $P = 0.019$), but when patients were dichotomized on the basis of PA, MSI remained inversely related to pre-PPCI mRPP in patients without PA ($r = -0.295$, $P = 0.004$) but not in those with PA ($r = -0.103$, $P = 0.422$) (Figure 3). Notably, in patients with PA, there was a positive relationship between MSI and the number of PA episodes (Spearman's $\rho = 0.332$, $P = 0.010$). On the other hand, no relationship was observed between MSI and the time elapsed from the first or last episode of PA to infarction ($r = 0.008$, $P = 0.949$ and $r = -0.133$, $P = 0.316$, respectively). At univariate linear regression

analysis, shorter time-to-PPCI, lower pre-PPCI mRPP, higher post-PPCI TIMI flow-grade, and the occurrence of PA were significantly associated with larger MSI (Table 4). At multivariate linear regression analysis, the occurrence of PA remained significantly associated with MSI after correction for pre- and post-PPCI TIMI flow-grade, pre-PPCI mRPP, and time-to-PPCI (Table 5).

Discussion

We showed that patients with acute MI experiencing new-onset PA in the week preceding the infarction had higher MSI, less infarct transmural, and lower incidence of MH and MO than patients without PA, resulting in better regional and global LV systolic function. Noteworthy, the occurrence of PA remained significantly associated to higher MSI even after correction for other factors implicated in ischaemic injury, including pre- and post-procedural TIMI flow-grade, myocardial oxygen consumption before PPCI, and time to revascularization. Moreover, in patients with PA, the degree of MSI was positively related to the number of episodes of PA.

Our study is novel in so far as we assessed the impact of new-onset PA on myocardial salvage using CMR in the early post-infarction period after accounting for factors known to influence the ischaemic damage and which could be readily assessed. In comparison to patients without PA, those with new-onset PA showed greater MSI, as a result of smaller MI size, less infarct transmural, and lower incidence of MH and MO. Previous studies have reported that coronary occlusion in the early period of infarction is highly dynamic,¹⁸ and Andreotti *et al.*⁵ have shown that in acute MI patients treated with thrombolysis, the occurrence of PA results in more rapid reperfusion, thus leading to smaller infarct size. However, in our study, the residual blood flow to the myocardium at risk through the infarct-related artery or epicardial collaterals was similar in patients with and without PA. Additionally, when the analysis was limited to patients with pre-procedural TIMI flow-grade ≤ 1 and Rentrop-grade ≤ 1 , the degree of MSI remained higher and infarct transmural was lower in patients with PA. These data indicate that IPC is likely a mechanism by which new-onset PA protects the myocardium at risk.³

Experimentally, the cardioprotective effect of IPC has been confirmed by a plethora of data derived from diverse animal species. In particular, it has been demonstrated that repetitive brief episodes of ischaemia protect the myocardium from the subsequent sustained ischaemia, limiting final MI size.^{3,19} Our findings expand previous observations made in humans that identified IPC as the putative mechanism underlying the beneficial effects of new-onset PA on post-infarction LV function recovery,^{4,6,7,9,10} as well as on short- and long-term prognosis.^{8,20,21} However, prior studies were limited by the lack of direct measure of salvaged myocardium and by their retrospective nature which did not allow to adjust for factors potentially influencing myocardial salvage. In our study, we prospectively analysed the variables known from previous experimental and clinical studies to intervene on ischaemic damage. Noteworthy, multivariate analysis confirmed that new-onset PA in the week preceding MI was independently associated with a higher MSI. Recently, in a subgroup analysis of a randomized controlled trial demonstrating the cardioprotective effect of exenatide




Figure 2 T2w-imaging (A and C) and LGE-imaging (B and D) in two patients with acute infero/lateral MI. One patient referred two episodes of PA within 48 h from infarction (upper panel), whereas the second patient was asymptomatic in the week preceding MI (bottom panel). Both patients presented the same angiographic characteristics (pre-/post-PPCI: 0/3; Rentrop-grade: 0) and similar time-to-PPCI (233 vs. 247 min) and mRPP before PPCI (10 143 vs. 10 127 [bpm] mmHg), nevertheless the patient with PA had higher MSI (0.68 vs. 0.05) and lower infarct transmuralty (34 vs. 98%) when compared with patient without PA. Additionally, MH (C, arrow) and MO (D, arrow) were detected only in the patient without PA.




Figure 3 Scatter plots and regression lines of MSI and logarithm (Log) of time-to-PPCI (A) and mRPP before PPCI (B) in the whole study population (black regression line) and in patients with (red triangles, red regression line) and without PA (blue circle, blue regression line).

Table 4 Univariate linear regression analysis for MSI determinants

Baseline characteristics	β -value	P-value
Age	0.010	0.901
Female gender	0.072	0.376
Hypertension	0.018	0.825
Diabetes	0.060	0.458
Hypercholesterolaemia	0.024	0.769
Family history of CAD	-0.050	0.543
Current smoking	0.014	0.861
Killip class	-0.029	0.726
Time-to-PPCI	-0.191	0.018
Prodromal angina	0.367	<0.001
Infarct-related artery	-0.012	0.883
TIMI flow-grade pre-PPCI	0.146	0.066
Presence of collaterals	-0.028	0.732
TIMI flow-grade post-PPCI	0.185	0.022
mRPP before PPCI	-0.191	0.019
Beta-blocker before PPCI	0.020	0.808
Nitrate before PPCI	-0.100	0.227
Morphine before PPCI	-0.017	0.841
Gp IIb/IIIa before PPCI	0.083	0.312

Abbreviations as reported in previous tables.

Table 5 Multivariate linear regression analysis for MSI determinants

Baseline characteristics	β -value	P-value
Time-to-PPCI	-0.172	0.018
Prodromal angina	0.352	<0.001
TIMI flow-grade pre-PPCI	0.114	0.121
TIMI flow-grade post-PPCI	0.180	0.013
mRPP before PPCI	-0.217	0.003

Abbreviations as reported in previous tables.

in acute MI patients, Lonborg *et al.*¹¹ showed that PA in the 3 months preceding MI was associated with higher MSI as quantified by CMR. However, this study had several limitations making the interpretation of the results difficult. First, it cannot be excluded that exenatide influenced the sub-study results; secondly, no data on MH and MO were reported; thirdly, the time interval between PA and infarction was not mentioned.

Experimentally, the cardioprotective effect of IPC is lost if the interval between the preconditioning stimulus and infarction exceeded few hours (classical IPC) to then reappear between 24 and 72 h after the initial IPC stimulus (delayed or 'second window' of IPC).^{3,19} In humans, classic IPC has been reported during percutaneous coronary angioplasty²² and cardiac surgery,²³ and the occurrence of new-onset PA within 24 h from acute MI has been associated with smaller MI size and

better post-infarction LV function recovery⁹ and prognosis.²⁰ Other groups have reported that PA within 48 h from acute MI is also associated with enhanced recovery of LV function after infarction^{4,8} and better in-hospital prognosis.⁷ In our study, all except two patients and more than half of patients experienced the last episode of PA within 48 and 6 h from infarction, respectively, indicating that classical or delayed IPC may account for the cardioprotective effect of PA. However, MSI was not related to the time elapsed from the first or last PA episode to acute MI. The independence of the time interval between episodes of PA and the degree of salvaged myocardium may be explained by the fact that silent ischaemia also contributes to IPC. Although one may argue that silent ischaemia might also occur in patients without PA, it has been reported that patients with PA had higher prevalence of silent pre-infarction ischaemia on 24 h Holter monitoring compared with asymptomatic patients.²⁴ Furthermore, we also found that in patients with PA, the degree of MSI was positively related to the number of new-onset PA episodes. Whether this finding indicates a cumulative cardioprotective effect of IPC in humans cannot be clarified by our results and merits further investigation. Reasonably, our findings indicate that patients with at least one episode of PA are more likely to be effectively preconditioned than asymptomatic patients, and the likelihood of cardioprotection is positively linked to the increasing number of PA episodes.

Interestingly, we also found that patients with PA had a lower incidence of MH and MO than patients without PA, indicating a more favourable pattern of post-infarction damage. Myocardial haemorrhage invariably denotes a severe damage of coronary microvasculature with leakage of red blood cells from injured vessels, and it occurs only in reperfused MI thereby representing a marker of post-reperfusion damage.^{12,25,26} One may speculate that IPC mitigates post-reperfusion damage in humans with a particular beneficial effect on the coronary microvasculature. This finding is consistent with experimental studies showing that IPC minimizes post-reperfusion injury¹⁹ not only acting directly on cardiomyocytes but also by preserving coronary microvasculature function.²⁷ Notably, in humans Komamura *et al.*²⁸ demonstrated that in patients with acute anterior MI, the occurrence of new-onset PA is associated with higher post-reperfusion great cardiac vein flow, reflecting better coronary microvascular function and smaller infarct size compared with patients without PA. Similarly, in acute MI patients undergoing PPCI, Takahashi *et al.*²⁹ reported that no-reflow phenomenon was less commonly observed in patients with PA compared with those without PA.

According to our study results, the assessment of new-onset PA in the week before infarction is of importance in studies testing novel or adjunctive reperfusion strategies aiming to protect the myocardium at risk, since the unbalanced distribution of this variable between the intervention and control groups may alter the final results.

Limitations

This was a three-centre study using different vendor CMR units, even though a similar study protocol was used with centralized data analysis. The determination of AAR by T2-weighted has limitations due to the inherently low signal-to-noise ratio and the susceptibility of signal loss in cardiac structures distant from the

surface coil. However, all CMR units used an SI correction algorithm to homogenize signal, and only patients with CMR exams showing a good or excellent T2-weighted images were included in the study. The parameters of T2-STIR sequence were slightly different across the diverse centres potentially resulting in discrepancies in the quantification of the AAR. We did not provide clinical nor CMR follow-up data. However, the study intended to investigate the pathophysiological factors influencing ischaemic damage in patients with acute MI. It is very likely that the beneficial effects of PA in the early post-infarction period will translate into better LV remodelling at mid- and long-term follow-up. Finally, the assessment of PA relied on the quality of history taking in acutely hospitalized patients.

Conclusion

New-onset PA in the week preceding acute MI is associated with larger myocardial salvage independent of other factors known to influence the jeopardized myocardium. The PA beneficial effect on salvaged myocardium is also associated with less coronary microvasculature damage.

Supplementary material

Supplementary material is available at *European Heart Journal – Cardiovascular Imaging* online.

Conflict of interest: none declared.

References

- Reimer KA, Jennings RB, Cobb FR, Murdock RH, Greenfield JC Jr, Becker LC et al. Animal models for protecting ischemic myocardium: results of the NHLBI Co-operative Study. Comparison of unconscious and conscious dog models. *Circ Res* 1985;**56**:651–5.
- Christian TF, Schwartz RS, Gibbons RJ. Determinants of infarct size in reperfusion therapy for acute myocardial infarction. *Circulation* 1992;**86**:81–90.
- Murry CE, Jennings RB, Reimer KA. Preconditioning with ischemia: a delay of lethal cell injury in ischemic myocardium. *Circulation* 1986;**74**:1124–36.
- Hirai T, Fujita M, Yamanishi K, Ohno A, Miwa K, Sasayama S. Significance of pre-infarction angina for preservation of left ventricular function in acute myocardial infarction. *Am Heart J* 1992;**124**:19–24.
- Andreotti F, Pasceri V, Hackett DR, Davies GJ, Haider AW, Maseri A. Preinfarction angina as a predictor of more rapid coronary thrombolysis in patients with acute myocardial infarction. *N Engl J Med* 1996;**334**:7–12.
- Nakagawa Y, Ito H, Kitakaze M, Kusuoka H, Hori M, Kuzuya T et al. Effect of angina pectoris on myocardial protection in patients with reperfused anterior wall myocardial infarction. *J Am Coll Cardiol* 1995;**25**:1076–83.
- Napoli C, Liguori A, Chiariello M, Di Ieso N, Condorelli M, Ambrosio G. New-onset angina preceding acute myocardial infarction is associated with improved contractile recovery after thrombolysis. *Eur Heart J* 1998;**19**:411–9.
- Kloner RA, Shook T, Przyklenk K, Davis VG, Junio L, Matthews RV et al. Previous angina alters in-hospital outcome in TIMI 4. *Circulation* 1995;**91**:37–45.
- Noda T, Minatoguchi S, Fujii K, Hori M, Ito T, Kanmatsuse K et al. Evidence for the delayed effect in human ischemic preconditioning: prospective multicenter study for preconditioning in acute myocardial infarction. *J Am Coll Cardiol* 1999;**34**:1966–74.
- Ottani F, Galvani M, Ferrini D, Sorbello F, Limonetti P, Pantoli D et al. Prodromal angina limits infarct size. A role for ischemic preconditioning. *Circulation* 1995;**91**:291–7.
- Lønborg J, Kelbæk H, Vejstrup N, Ahtarovski KA, Kelbæk H, Andersen M et al. Influence of pre-infarction angina, collateral flow, and pre-procedural TIMI flow on myocardial salvage index by cardiac magnetic resonance in patients with ST-segment elevation myocardial infarction. *Eur Heart J Cardiovasc Imaging* 2012;**13**:433–43.
- Ganame J, Messalli G, Dymarkowski S, Rademakers FE, Desmet W, Van de Werf F et al. Impact of myocardial haemorrhage on left ventricular function and remodelling in patients with reperfused acute myocardial infarction. *Eur Heart J* 2009;**30**:1440–9.
- Masci PG, Ganame J, Francione M, Desmet W, Lorenzoni V, Iacucci I et al. Relationship between location and size of myocardial infarction and their reciprocal influences on post-infarction left ventricular remodelling. *Eur Heart J* 2011;**32**:1640–8.
- Gobel FL, Norstrom LA, Nelson RR, Jorgensen CR, Wang Y. The rate-pressure product as an index of myocardial oxygen consumption during exercise in patients with angina pectoris. *Circulation* 1978;**57**:549–56.
- Rentrop KP, Cohen M, Blanke H, Philips RA. Changes in collateral channel filling immediately after controlled coronary artery occlusion by an angioplasty balloon in human subjects. *J Am Coll Cardiol* 1985;**5**:587–92.
- Bondarenko O, Beek AM, Hofman MB, Kühl HP, Twisk JW, van Dockum WG et al. Standardizing the definition of hyperenhancement in the quantitative assessment of infarct size and myocardial viability using delayed contrast enhanced CMR. *J Cardiovasc Magn Reson* 2005;**7**:481–5.
- Cerqueira MD, Weissman NJ, Dilsizian V, Jacobs AK, Kaul S, Laskey WK et al. AHA Writing Group on Myocardial Segmentation and Registration for Cardiac Imaging. *Circulation* 2002;**105**:539–42.
- Hackett D, Davies G, Chierchia S, Maseri A. Intermittent coronary occlusion in acute myocardial infarction. *N Engl J Med* 1987;**317**:1055–9.
- Vinten-Johansen J, Zhao AQ, Zatta AJ, Dobson GP. Preconditioning and postconditioning: innate cardioprotection from ischemia-reperfusion injury. *J Appl Physiol* 2007;**103**:1441–8.
- Ishihara M, Sato H, Tateishi H, Kawagoe T, Shimatani Y, Kurisu S et al. Implication of prodromal angina pectoris in anterior wall infarction. *J Am Coll Cardiol* 1997;**30**:970–5.
- Bahr RD, Leino EV, Christenson RH. Prodromal unstable angina in acute myocardial infarction: prognostic value of short- and long-term outcome and predictor of infarct size. *Am Heart J* 2000;**140**:126–33.
- Tomai F, Crea F, Chiarello L, Gioffre PA. Ischemic preconditioning in humans: models, mediators and clinical relevance. *Circulation* 2001;**100**:559–63.
- Yellon DM, Alkhalafi AM, Pugsley WB. Preconditioning the human myocardium. *Lancet* 1993;**342**:276–7.
- Nademanee K, Intarachot V, Josephson MA, Rieders D, Vaghaiwalla Mody F, Singh BN. Prognostic significance of silent myocardial ischemia in patients with unstable angina. *J Am Coll Cardiol* 1987;**10**:1–9.
- Fishbein MC, Y-Rit J, Lando U, Kanmatsuse K, Mercier JC, Ganz W. The relationship of vascular injury and myocardial haemorrhage to necrosis to reperfusion. *Circulation* 1980;**6**:1274–9.
- Garcia-Dorado D, Théroux P, Solares J, Alonso J, Fernandez-Avilés F, Elizaga J et al. Determinants of hemorrhagic infarcts. Histologic observations from experiments involving coronary occlusion, coronary reperfusion, and reocclusion. *Am J Pathol* 1990;**137**:301–11.
- Kaeffer N, Richard V, Thuillez C. Delayed coronary endothelial protection 24 hours after preconditioning: role of free radicals. *Circulation* 1997;**96**:2311–6.
- Komamura K, Kitakaze M, Nishida K, Naka M, Tamai J, Uematsu M et al. Progressive decreases in coronary vein flow during reperfusion in acute myocardial infarction. *J Am Coll Cardiol* 1994;**24**:370–7.
- Takahashi T, Anzai T, Yoshikawa T, Asakura Y, Ishikawa S, Mitamura H et al. Absence of preinfarction angina is associated with a risk of no-reflow phenomenon after primary coronary angioplasty for a first anterior wall acute myocardial infarction. *Int J Cardiol* 2000;**75**:253–60.